

The Ecology of Episodic Saline Lakes of Inland Eastern Australia, as Exemplified by a Ten Year Study of the Rockwell-Wombah Lakes of the Paroo.

BRIAN V. TIMMS

School of Environmental and Life Sciences, University of Newcastle, Callaghan, NSW, 2308, Australia.
Email: brian.timms@newcastle.edu.au

Timms, B.V. (2008). The ecology of episodic saline lakes of inland eastern Australia, as exemplified by a ten year study of the Rockwell-Wombah Lakes of the Paroo. *Proceedings of the Linnean Society of New South Wales* **129**, 1-16.

Studies on salt lakes are mostly snapshots of their unique characteristics and relationships. Longer term studies provide different perspectives on variability in hydrology, salinity and biological communities. Such a study on five lakes near the Paroo River in the northwestern Murray-Darling Basin showed most hold water episodically for about 80% of the time, but each fluctuate over a characteristic salinity range : unnamed lake 0.6 – 19 gL⁻¹, Wombah 1.2 – 30 gL⁻¹, North Blue 0.3 – 31 gL⁻¹, Mid Blue 0.7 – 103 gL⁻¹, and Bulla 1.8 – 262 gL⁻¹. Generally, instantaneous biodiversity is low and not necessarily correlated with salinity, but unlike southern seasonal salt lakes, species accumulation lists are long, approaching 80 species of invertebrates, 50 bird species and a few fish species per lake. Diversity is promoted by salinity fluctuation and habitat heterogeneity. Most species reach peak abundance in any season as long as conditions are within their physiological salinity tolerances. Invertebrate fauna is of inland affinities, but with some localized substractions and additions explained by hydrology and/or salinity; waterbird numbers are influenced by events elsewhere in Australia as well as by local conditions. Like most naturally salinised lakes, production can be high, especially at low to moderate salinities and algal blooms occur naturally from time to time.

Manuscript received 18 April 2007, accepted for publication 19 September 2007.

KEYWORDS: benthos, biodiversity, episodic lakes, fish, littoral, salinity variability, waterbirds, zooplankton.

INTRODUCTION

Over the last few decades much has been learnt about the numerous salt lakes in Australia. Limnological summaries are available for Victoria (Williams 1981) and the eastern inland (i.e. Queensland, New South Wales, adjacent South Australia) (Timms 2007) and southwestern Western Australia (Pinder et al. 2004a), Tasmania (De Deckker and Williams 1982), and southern South Australian lakes (De Deckker and Geddes 1982; Williams 1984). In essence, most Australian salt lakes are shallow, intermittent (either seasonal or episodic), chemically dominated by NaCl, markedly alkaline, and have a crustacean-dominated fauna that decreases in diversity with increasing salinity (see above references). Additionally, unlike salt lakes in the northern hemisphere, there is considerable regionalisation of the fauna (Williams 1984; Timms 2007).

Australian saline lakes are changing due to

various adverse environmental pressures, as reviewed by Williams (2002) and Timms (2005). In Australia the greatest problem is secondary salinisation and it is manifest mainly in southwestern Western Australia (Davis et al. 2003; Halse et al. 2003) and to a lesser degree in some areas of southern Victoria and South Australia (e.g. Lake Baird, Pellana Lagoon, Lake Wangary on Eyre Peninsula (author, unpublished data))(Fig 1 in Timms, 2005). However in the remote inland, saline systems have remained unaffected by anthropogenic secondary salinisation.

One such inland area is the middle and lower Paroo River catchment in northwestern New South Wales and adjacent southwestern Queensland (Fig. 1 in Timms, 2006). Here there are many freshwater wetlands but a few are naturally salinised. Although surrounding terrestrial environments are degraded after 120 years of grazing, aquatic habitats are almost pristine (Timms 2001a). Study of them provides a 'control' for investigations into secondary salinised

ECOLOGY OF EPISODIC SALINE LAKES



Figure 1. Map showing the five lakes on Rockwell and Wombah stations, southwest Queensland.

systems. However, they have accumulated their salt over millennia, so the rate of salinisation is different to that in anthropogenically affected systems. Also, the latter are affected by multiple degrading factors rather than just by salt increase (Davis et al. 2003).

Much of the existing data on saline lakes were collected on single field trips, or perhaps at best, a few over a year, so that long-term fluctuations associated with variable weather patterns have gone unrecognised. Yet these fluctuations could be a major difference between primary and secondary salinised lakes (Hudson et al. 2003). So understanding of inland lakes over a longer time frame than provided by most studies will provide better understanding of some of the impacts in secondary salinised systems. The present study aims to do this in a context of other saline lakes in the Paroo and of Australia in general.

THE LAKES

The five lakes lie on a creek line (called Number 10 creek upstream of North Blue Lake) that flows southwestward to join an anabranch of the Paroo River on the New South Wales – Queensland border 40-50 km east of Hungerford (Fig. 1). The creek is channelled between the upper three lakes (North Blue, Mid Blue and Bulla) which coalesce when full, but is obliterated by dunes downstream of Lake Bulla. It reappears as a small stream near Lake Wombah (Fig. 1). Paroo floodwater often enters Lake Wombah and leaves by the same route, but rarely reaches upstream to Lake Bulla. The last times this occurred were in 1974 and 1990 (P. Tuite, pers com.). The upper three lakes fill from Number 10 Creek, together with local streams from the east. The unnamed lake fills entirely

Table 1. Geomorphic and physicochemical features of the five lakes

| Parameter | Lake Wombah | Unnamed Lake | Lake Bulla | Mid Blue Lake | Nth Blue Lake |
|--------------------------------|-------------|--------------|------------|---------------|---------------|
| Area (ha) | 740 | 70 | 420 | 210 | 205 |
| Depth (m)* | 2.3 (3.2) | 0.2 (1.8) | 4.4 (4.9) | 3.3 (3.9) | 2.3 (3.6) |
| % of years with water | 80 | 15 | 82 | 82 | 75 |
| TDS range mgL ⁻¹ | 1.2 - 30 | 0.6 - 19 | 1.8 - 262 | 0.7 - 103 | 0.3- 31 |
| Mean TDS (gL ⁻¹) | 7.6 | 5.6 | 34 | 9.1 | 6.8 |
| Median TDS (gL ⁻¹) | 4.9 | 3.3 | 9.8 | 4.1 | 4.2 |
| Mean water Temperature (° C) | 22 | 19.5 | 19.4 | 23.6 | 23.8 |
| Mean Turbidity ± SE (FTU) | 34 ± 10.9 | 12 ± 2.7 | 7 ± 1.7 | 14 ± 3.5 | 13 ± 5.1 |
| Mean pH ± SE | 9 ± 0.21 | 8.7 ± 0.16 | 8.8 ± 0.19 | 8.9 ± 0.12 | 8.8 ± 0.23 |

* Figures in brackets are heights of highest water marks, but were not recorded during study.

from local streams and Lake Wombah receives most of its water from the Paroo River.

METHODS

Areas of the lakes were established from aerial photographs and depths from gauges in the lakes and the height of stranded beaches/wave cut notches determined with a dumpy level (Table 1). The later relate to intense local rainfall and large Paroo floods such as occurred in 1974 and 1990 (P. Tuite, pers. com.).

The lakes were visited at approximately three-monthly intervals from July 1995 to June 2004 for the purposes of determining some physicochemical parameters and sampling zooplankton and littoral organisms. On each visit a surface water sample from about 50m offshore was taken for the immediate measurement of temperature with a mercury thermometer, pH with a HANNA HI8924 meter and later measurement of total dissolved solids (TDS) by gravimetry, with turbid waters being allowed to settle for many months in a sealed container. Turbidity was measured spectrophotometrically in the laboratory at 450 nm with the results recorded in Fittou's Turbidity

Units. Five times during 1997 and 1998 nitrate and phosphate was determined in the field on the water samples from Wombah, Bulla, Mid Blue and North Blue using a HACH DR/2000 spectrophotometer and method 8171 for dissolved nitrate and 8048 for dissolved phosphate. These measurements were made during a hyposaline and a meso/hypersaline stage. Nutrients were measured once only (July 1998) in the unnamed lake.

Zooplankton and littoral organisms were collected with nets of mesh size 159 µm and 1mm respectively, identified and counted as outlined in Timms and McDougall (2004). Benthos was sampled with a Birge-Ekman grab just once in the four larger lakes— Wombah in April 2004 and Bulla, Mid Blue and North Blue in December 2001. Five grabs were taken from each lake near the deepest part, sieved onshore through a 0.4 mm mesh and sorted while organisms were alive. Fish were caught sometimes in the littoral net and often in yabbie traps set for an hour or two to catch *Cherax* crayfish. These traps were not employed regularly or in any pattern.

Waterbirds were counted from January 1998 onwards. Shore based total counts were made with binoculars and/or a spotting scope on each visit. Counts were always made from the same vantage

ECOLOGY OF EPISODIC SALINE LAKES

points each time, and covered the full surface area of all lakes except Lake Wombah. In this lake only the northern half was surveyed, for if the southern part was included it would take several hours to count the birds and impossible to account for bird movements because of the terrain. Small waders were underestimated in Lakes Wombah and Bulla when full as distances were too great for accurate determination.

RESULTS

Physicochemical Conditions

Average annual rainfall for the 10-year period was 366 mm, but yearly totals varied widely and included unusually wet years of 1998 (618 mm) and 2000 (622 mm) and the particularly dry year of 2002 (92 mm) (Fig. 2). Major local inflows to the top three lakes from intense local rain occurred in April 1998 and March-May 2000, and Paroo floods filled Lake Wombah in July 1998, March 2000 and February 2004. Between major fillings lake levels fell slowly with evaporation or rose slightly with minor inputs, so that four of the lakes held water for ca 80% of the time (Table 1). High salinities were experienced only briefly during March to December 1997 (to April 1998 in Bulla) and again variously during much of 2001-2002 in Wombah, Bulla, Mid Blue and North Blue (Fig. 3). The small unnamed lake held water for

a few months in 1998-99 and again in 2000, both as a consequence of unusually heavy rainfall in those years (see above).

All five lakes varied widely in salinity, from fresh in all cases to hyposaline (i.e. to ca 20gL^{-1}) in the unnamed lake, to mesosaline (i.e. $20\text{-}50\text{gL}^{-1}$) in Wombah and North Blue, to hypersaline (i.e. $>50\text{gL}^{-1}$) in Mid Blue and very hypersaline ($>200\text{gL}^{-1}$) in Bulla (Table 1). Mean salinities are deceptive due to the short time the lakes spend at higher salinities (Fig 3) so that median salinities are more representative than mean salinities of typical conditions in the lakes (Table 1); in this respect all lakes are often hyposaline, with Bulla the most saline and the unnamed lake the least.

Mean water temperature for the ten years was $21.6\text{ }^{\circ}\text{C}$, with a minimum of $8\text{ }^{\circ}\text{C}$ and maximum of $36\text{ }^{\circ}\text{C}$. Lake waters were generally clear, except following major inflows, or when shallow and wind-stirred, or in Lake Wombah's river-derived water (Table 1). Variability was greatest in the two lakes (Wombah and North Blue) receiving mostly flood water and least in Bulla where abundant salt helped to settle colloidal clay (Table 1). All five lakes had well buffered, markedly alkaline water, except again for some variation in the two lakes receiving most floodwater (Table 1).

Nutrients were high, around $1\text{-}2\text{ mgL}^{-1}$ nitrate and $0.2\text{-}0.4\text{ mgL}^{-1}$ phosphate, with highest values in Lake Bulla (Table 1).




Figure 3. Fluctuations in TDS in the five lakes.




Figure 2. Monthly rainfall at Rockwell station homestead 1996-2004.

ECOLOGY OF EPISODIC SALINE LAKES

Aquatic Plants

Aquatic macrophytes were most common and persistent in the clear waters of Lake Bulla, where *Myriophyllum verrucosum* Lindl, *Lepilaena bilocularis* Kirk, and *Chara* spp. were common at lower salinities (<30 g/L). At higher salinities (30-60 g/L) the dominants were *Chara* sp. and *Ruppia* sp. These species occurred in the other lakes, but were less abundant, except occasionally in North Blue. A new species of *Chara* occurs in Bulla and Mid Blue (A. Garcia, pers. comm.) In summer and autumn, filamentous algae often shrouded plants, and appeared to be associated with their demise for the season.

Zooplankton

At least 37 taxa occur in the five lakes, with 12-28 species per lake (Table 2). The dominant freshwater species were *Boeckella triarticulata*, *Daphnia angulata* and *D. lumholtzi*, while the most common saline species were *Apocyclops dengizicus*, *Daphniopsis queenslandensis*, *Cyprinotus* sp. and *Diacypris* spp. (Table 2). There are broad similarities between average percentage composition between the lakes (Table 2), though the most intermittently filled unnamed lake had the highest percentage of clam shrimps (14.3%) and the most saline lake (Bulla) had the highest percentage of saline species (69.9%), followed in order by Mid Blue (56.0%), North Blue (44.4%) and Wombah (14.7%). Freshwater eulimnetic species with limited salinity tolerance (e.g. *Calamoecia lucasi*, *Daphnia lumholtzi*, *Daphnia angulata*, *Ceriodaphnia* spp., *Diaphanosoma* spp., *Moina australiensis*) comprised <22% in all lakes and were lowest in Bulla (10.2%) and the unnamed lake (1.1%). While salinity influenced a few species, momentary species richness was not related significantly to salinity in any of the lakes (Wombah $r = -0.2148$, $n=27$; unnamed lake $r = 0.0832$, $n=7$; Bulla $r = -0.4885$, $n=31$; Mid Blue $r = -0.1425$, $n=35$; North Blue $r = -0.2505$, $n=29$).

Typically, composition and dominance varied markedly between sampling dates in each lake and there was little seasonal repeatability over the ten years. Exceptions to this were *Daphnia lumholtzi* which often peaked in autumn in the fresher lakes, and *Daphniopsis queenslandensis* which was most common in winter-spring and *Moina baylyi* in summer-autumn, when lakes were saline. The most predictable occurrences were of saline species when lake salinity was elevated (e.g. *Daphnia* n.sp., *Cyprinotus* sp. at lower salinities, and *Apocyclops dengizicus* and *Diacypris* spp. at higher salinities).

Table 2 (RIGHT) . Zooplankton of the five lakes, showing mean percentage composition and maximum salinity for each species.

The amount of zooplankton in each sample varied greatly, with means ranging from 22.2 (Bulla) to 4.1 (North Blue) mls per minute of standardised collecting (Table 3). Variability was greatest in the more saline lakes (Table 3). Dense populations were relatively uncommon but sometimes occurred in winter, soon after filling, or when the lakes increased in salinity as they dried rapidly (as a consequence of long periods of warm dry weather), but this was not significant statistically.

Table 3. Measure of standing crops

| Lake | Zooplankton | Birds* | Benthos |
|----------|-----------------------|-----------------|----------------------------|
| | volume \pm SE (mls) | no./ha \pm SE | g m ⁻² \pm SE |
| Wombah | 19.4 \pm 5.4 | 7.3 \pm 2.3 | 9.1 \pm 1.1 |
| unnamed | 16.6 \pm 2.6 | | |
| Bulla | 22.2 \pm 8.1 | 15.5 \pm 5.5 | 33.5 \pm 0.9 |
| Mid Blue | 9.4 \pm 2.9 | 13.2 \pm 4.3 | 4.5 \pm 0.7 |
| Nth Blue | 4.1 \pm 0.8 | 18.6 \pm 3.6 | 4.9 \pm 0.4 |

*densities based on lakes being on average half full

Littoral Invertebrates

At least 78 species occur in the five lakes, with an accumulated species richness in each lake between 36-58 species (Table 4). Mean momentary species richness in each lake ranged from 6.4 to 10.9 (Table 4). The dominant species in all lakes was *Micronecta* sp., with the two ostracods *Mytilocypris splendida* and *Trigonocypris globulosus* and *Anisops gratus* and other backswimmers and boatmen common in most lakes, except in the unnamed lake. All common species had wide salinity tolerances (Table 4). Though *Tanytarsus barbitarsus* had by far the highest salinity tolerance, it was common only in mesosaline waters and above.

Beetles, although diverse, were not common in the lakes, and larvae were uncommon too. Odonatans were most abundant in lakes with extensive macrophytes (Bulla, Mid Blue). Ephemeropterans were most abundant in the lower salinity lakes (unnamed, Mid Blue and North Blue). River dominants such as large crustaceans *Macrobrachium australiense* and *Cherax destructor* were hardly encountered and then only in

B.V. TIMMS

| SPECIES | max. salinity | Lake Wombah | unnamed Lake | Bulla Lake | Mid Blue Lake | Nth Blue Lake |
|--|------------------|-------------|--------------|------------|---------------|---------------|
| | gL ⁻¹ | n=27 | n=7 | n=31 | n=35 | n=29 |
| Spinicaudata | | | | | | |
| <i>Caenestheria</i> n.sp. | 3.3 | | 13.6 | | | |
| <i>Caenestheriella packardi</i> Spencer & Hall | 0.7 | | | | <0.1 | <0.1 |
| <i>Eocyclus parooensis</i> Richter & Timms | 3.3 | <0.1 | 0.7 | | | |
| Copepoda | | | | | | |
| <i>Boeckella triarticulata</i> Thomson | 19 | 48.0 | 54.5 | 18.5 | 12.8 | 16.9 |
| <i>Calamoecia lucasi</i> Brady | 6.7 | 4.0 | | 1.5 | 0.3 | 2.1 |
| <i>Thermocyclops decipiens</i> Kiefer | 7.7 | 7.8 | 15.7 | | 1.2 | 8.0 |
| <i>Mesocyclops kieferi</i> Van de Velde | 4.2 | 4.9 | | | 4.2 | 5.3 |
| <i>Apocyclops dengizicus</i> (Lepeschkin) | 94 | 4.4 | | 30.2 | 24.3 | 13.9 |
| <i>Cletocamptus deitersi</i> (Richard) | 13 | 0.1 | | 0.5 | 0.9 | 0.1 |
| Cladocera | | | | | | |
| <i>Daphnia lumholtzi</i> Sars | 6.7 | 0.1 | | 2.8 | 9.4 | 6.9 |
| <i>Daphnia angulata</i> Hebert | 5.4 | 7.9 | 0.9 | 2.4 | 7.9 | 9.4 |
| <i>Daphnia cephalata</i> King | 2.1 | | | 0.1 | | |
| <i>Daphnia</i> n. sp. | 16 | | | 3.4 | 0.5 | 2.3 |
| <i>Daphniopsis queenslandensis</i> Sergeev | 59 | 1.9 | | 9.5 | 9.5 | 8.8 |
| <i>Ceriodaphnia cornuta</i> Sars | 2.1 | | | | <0.1 | |
| <i>Ceriodaphnia</i> n.sp. | 1.9 | <0.1 | 0.2 | 1.6 | 2.6 | 2.8 |
| <i>Simocephalus vetulus</i> (King) | 0.7 | | | | | <0.1 |
| <i>Diaphanosoma excisum</i> | 3.1 | 1.8 | | <0.1 | | |
| <i>Diaphanosoma unguiculatum</i> Gurney | 1.7 | | | | | 1.2 |
| <i>Moina australiensis</i> Sars | 4.9 | 4.9 | | | | |
| <i>Moina baylyi</i> Forro | 59 | 0.1 | | 3.1 | 0.7 | 0.3 |
| <i>Macrothrix carinata</i> (Smirnov) | 8.5 | 0.1 | 0.6 | 0.2 | 1.3 | 0.2 |
| <i>Latonopsis australis</i> Sars | 4.2 | | | | 0.2 | |
| <i>Alona</i> spp. | 11 | 4.6 | | 0.2 | 1.1 | 2.0 |
| <i>Celsinotum</i> spp. | 19 | 0.2 | 2.4 | 0.4 | 0.7 | 0.2 |
| <i>Dunhevedia</i> sp. | 11 | | | 0.3 | 0.4 | <0.1 |
| <i>Leydigia</i> sp. | 5.9 | | | | | 0.4 |
| Ostracoda | | | | | | |
| <i>Bennelongia</i> sp. | 6.2 | 0.9 | 1.9 | | 1.7 | |
| <i>Cyprinotus</i> sp. | 16 | 3.0 | 8.4 | 9.0 | 9.1 | 9.1 |
| <i>Diacypriis</i> spp. | 157 | 2.1 | | 14.1 | 8.3 | 2.5 |
| <i>Heterocypris</i> sp. | 41 | 0.3 | | 0.9 | 0.5 | 4.5 |
| <i>Ilyocypris</i> sp. | 2.2 | | | | <0.1 | <0.1 |
| <i>Newnhamia</i> sp. | 2.2 | | | | <0.1 | <0.1 |
| <i>Reticypris</i> sp. | 35 | 0.7 | | 0.2 | 1.7 | 2.5 |
| <i>Mytilocypris splendida</i> (Chapman) | 40 | 1.8 | | 0.7 | 0.3 | 0.2 |
| <i>Trigonocypris globulosa</i> De Deckker | 46 | 0.2 | 1.0 | 0.4 | 0.4 | 0.1 |
| number of species | | 24 | 12 | 23 | 28 | 28 |

ECOLOGY OF EPISODIC SALINE LAKES

Table 4. Littoral invertebrates in the five lakes. All numbers expressed as mean log abundance (Log 1 = 10) individuals per 15 minute collection).

| | max. | Lake | unnamed | Bulla | Mid Blue | Nth Blue |
|--|----------|--------|---------|-------|----------|----------|
| Species | salinity | Wombah | Lake | Lake | Lake | Lake |
| | g/L-1 | n=23 | n=7 | n=28 | n=34 | n=26 |
| Anostraca | | | | | | |
| <i>Branchinella australiensis</i> (Richters) | 1 | | | | | <0.01 |
| <i>Branchinella buchananensis</i> Geddes | 3 | | 0.14 | | | |
| Spinicaudata | | | | | | |
| <i>Caenestheria</i> sp. | 3 | | 0.71 | | | |
| <i>Caenestheriella packardi</i> Spencer & Hall | 1 | 0.04 | 0.57 | <0.01 | 0.03 | 0.05 |
| <i>Eocyclus parooensis</i> Richter & Timms | 3 | | 0.86 | <0.01 | | |
| <i>Eocyclus</i> n sp | 1 | 0.06 | | | | |
| <i>Limnadopsis birchii</i> (Baird) | 3 | <0.01 | 0.14 | | | |
| Ostracoda | | | | | | |
| <i>Mytilocypris splendida</i> (Chapman) | 68 | 1.15 | | 1.36 | 0.74 | 0.62 |
| <i>Trigonocypris globulosa</i> De Deckker | 68 | 0.52 | 0.01 | 1.68 | 0.92 | 1.16 |
| Decapoda | | | | | | |
| <i>Cherax destructor</i> Clark | 4 | <0.01 | | | | |
| <i>Macrobranchium australiense</i> Riek | 6 | <0.01 | | | | |
| Ephemeroptera | | | | | | |
| <i>Cloeon</i> sp. | 5 | 0.09 | 0.31 | 0.07 | 0.30 | 0.35 |
| <i>Tasmanocoenis tillyardi</i> (Lestage) | 5 | | | | 0.30 | 0.16 |
| Odonata | | | | | | |
| <i>Austrolestes annulosus</i> (Selys) | 4 | | | 0.03 | <0.01 | |
| <i>Ischnura heterostricta</i> (Burmeister) | 30 | 0.05 | | 0.22 | 0.15 | 0.28 |
| <i>Xanthoagrion erythroneurum</i> Selys | 12 | 0.04 | 0.14 | 0.28 | 0.21 | 0.06 |
| <i>Diplacoides bipunctata</i> (Brauer) | 12 | | | 0.11 | 0.06 | <0.01 |
| <i>Hemicordulia tau</i> (Selys) | 12 | | 0.16 | 0.22 | 0.30 | 0.06 |
| <i>Hemianax papuensis</i> (Burmeister) | 3 | 0.04 | | 0.18 | 0.06 | |
| <i>Trapezostigma loweii</i> (Kaup) | 4 | | | 0.03 | | |
| Hemiptera | | | | | | |
| <i>Anisops calcaratus</i> Hale | 8 | 0.18 | <0.01 | 0.14 | 0.09 | <0.01 |
| <i>Anisops gratus</i> Hale | 26 | 1.2 | 1.01 | 0.53 | 0.71 | 0.35 |
| <i>Anisops stahi</i> Kirkaldy | 13 | | | | 0.03 | |
| <i>Anisops thienemanni</i> Lundbald | 12 | 0.35 | 0.59 | 0.18 | 0.43 | 0.36 |
| <i>Micronecta</i> sp. | 94 | 1.93 | 2.71 | 1.26 | 2.16 | 2.26 |
| <i>Agraptocorixa eurynome</i> Kirkaldy | 17 | 0.48 | 1.44 | 0.22 | 0.33 | 0.23 |
| <i>Agraptocorixa hirtifrons</i> Hale | 17 | 0.32 | 0.59 | 0.14 | 0.11 | 0.24 |
| <i>Agraptocorixa parvipunctata</i> Hale | 17 | 0.31 | | 0.07 | <0.01 | 0.04 |
| <i>Sigara</i> sp. | 3 | | | | 0.06 | |
| Trichoptera | | | | | | |
| <i>Notolina</i> sp. | 5 | | | | 0.06 | <0.01 |
| <i>Oecetis</i> sp. a | 5 | <0.01 | 0.14 | 0.04 | 0.03 | 0.05 |
| <i>Oecetis</i> sp. b | 5 | | | 0.03 | 0.06 | 0.04 |
| <i>Triplectides australicus</i> Banks | 11 | 0.17 | 0.44 | 0.11 | 0.33 | 0.20 |

B.V. TIMMS

| | | | | | | |
|---|-----|-------|-------|-------|-------|-------|
| Coleoptera | | | | | | |
| <i>Allodessus bistrigatus</i> (Clark) | 12 | 0.05 | 0.16 | 0.19 | 0.25 | 0.20 |
| <i>Antiporus gilberti</i> Clark | 8 | 0.08 | 0.43 | 0.18 | 0.21 | <0.01 |
| <i>Bidessoides</i> sp. | 8 | | | | 0.06 | |
| <i>Cybister tripuncatus</i> Olivier | 68 | 0.08 | | <0.01 | 0.04 | <0.01 |
| <i>Eretes australis</i> (Erichson) | 8 | 0.05 | | <0.01 | | |
| <i>Hydroglyphus leai</i> (Guignot) | 4 | | | | <0.01 | <0.01 |
| <i>Hyphydrus elegans</i> (Montrouzier) | 1 | <0.01 | | <0.01 | | <0.01 |
| <i>Megaporus howitti</i> (Clark) | 2 | | | | 0.03 | |
| <i>Platynectes decempunctatus</i> (Fabricius) | 2 | <0.01 | | | 0.03 | |
| <i>Rhantus suturalis</i> (MacLeay) | 8 | | <0.01 | <0.01 | 0.03 | |
| <i>Sternopriscus multimaculatus</i> (Sharp) | 12 | <0.01 | | <0.01 | 0.04 | |
| <i>Haliphus</i> sp. | 3 | | <0.01 | | 0.04 | 0.04 |
| <i>Hydraena</i> sp. | 3 | | 0.14 | | | |
| <i>Hydrochus</i> sp. | 1 | <0.01 | | | | |
| <i>Berosus approximans</i> Fairmaire | 1 | | | | 0.03 | 0.08 |
| <i>Berosus australiae</i> Mulsant | 2 | | 0.14 | | 0.06 | |
| <i>Berosus debillipennis</i> Blackburn | 3 | | | | <0.01 | |
| <i>Berosus macumbensis</i> Blackburn | 2 | | 0.29 | 0.04 | 0.03 | 0.04 |
| <i>Berosus munitipennis</i> Blackburn | 26 | 0.14 | 0.57 | 0.04 | 0.12 | <0.01 |
| <i>Berosus nutans</i> MacLeay | 12 | | 0.3 | | <0.01 | |
| <i>Enochrus andersoni</i> Blackburn | 7 | 0.06 | 0.29 | 0.11 | 0.12 | |
| <i>Enochrus maculiceps</i> (MacLeay) | 2 | | | | 0.03 | |
| <i>Hydrophilus brevispina</i> Fairmaire | 7 | | <0.01 | <0.01 | | <0.01 |
| <i>Limnoxenus zealandicus</i> (Broun) | 8 | | <0.01 | 0.04 | 0.06 | |
| Beetle larvae <i>Antiporus</i> sp. | | <0.01 | | 0.08 | <0.01 | |
| Beetle larvae <i>Eretes</i> sp. | | | <0.01 | | | |
| Beetle larvae <i>Rhantus</i> sp. | | | 0.43 | | 0.03 | |
| Beetle larvae <i>Berosus</i> sp. | | 0.09 | | 0.04 | 0.04 | <0.01 |
| Beetle larvae <i>Hydrophilus</i> sp. | | | 0.14 | | | <0.01 |
| Diptera | | | | | | |
| <i>Chironomus</i> spp. | 11 | 0.18 | 1.14 | 0.33 | 0.46 | 0.58 |
| <i>Coelopynia</i> sp. | 8 | 0.02 | | | | 0.01 |
| <i>Cryptochironomus</i> sp. | 8 | 0.21 | | <0.01 | 0.09 | 0.15 |
| <i>Dicrotendipes</i> sp. | 30 | 0.17 | 0.44 | 0.11 | 0.09 | 0.12 |
| <i>Polypedilum nubifer</i> Skuse | 17 | 0.26 | | 0.25 | 0.41 | 0.35 |
| <i>Procladius</i> sp. | 68 | 0.17 | | <0.01 | | 0.15 |
| <i>Tanytarsus barbitarsus</i> Freeman | 193 | | 0.04 | 0.54 | | 0.12 |
| unidentified ceratopogonid larvae | 34 | 0.04 | | 0.18 | 0.12 | 0.08 |
| <i>Aedes</i> sp. | 3 | | 0.43 | | | |
| <i>Anopheles</i> sp. | 12 | | 0.43 | | 0.06 | 0.04 |
| unidentified syrphid larvae | 1 | <0.01 | | | | <0.01 |
| unidentified tabanid larvae | 7 | | 0.29 | 0.22 | <0.01 | |
| Lepidoptera | | | | | | |
| unidentified pyralidae larvae | 4 | <0.01 | | 0.04 | 0.12 | 0.04 |

ECOLOGY OF EPISODIC SALINE LAKES

| Table 4 continued | | | | | | |
|----------------------------------|---------------|-------------|--------------|------------|---------------|---------------|
| Species | max. salinity | Lake Wombah | unnamed lake | Bulla Lake | Mid Blue Lake | Nth Blue Lake |
| Hydrocarina | | | | | | |
| <i>Arrenurus</i> sp. | 7 | <0.01 | | 0.18 | 0.13 | 0.12 |
| <i>Eylais</i> sp. | 5 | 0.22 | 0.31 | 0.04 | 0.18 | 0.04 |
| <i>Hydrachna</i> sp. | 8 | 0.13 | | | 0.06 | 0.16 |
| <i>Limnesia</i> sp. | 7 | 0.17 | | 0.14 | 0.21 | 0.42 |
| <i>Poina</i> sp. | 1 | | | <0.01 | | 0.04 |
| unidentified water mite | 2 | | | | 0.03 | 0.12 |
| Gastropoda | | | | | | |
| <i>Coxiella gilesi</i> (Angas) | 34 | | | 0.79 | 0.30 | |
| <i>Glyptophysa</i> sp. | 3 | | | 0.04 | 0.09 | 0.04 |
| momentary species richness (MSR) | | 6.4 | 10.9 | 7.1 | 9.5 | 8.0 |
| MSR SE | | 0.6 | 2.1 | 0.8 | 0.8 | 0.7 |
| number of species | | 45 | 36 | 49 | 58 | 51 |

Lake Wombah. *Coxiella gilesi* was abundant only in Lake Bulla and Mid Blue Lake. Chironomids were present in most lakes and were no doubt more diverse than indicated in Table 4. The unnamed lake was the most distinctive of the five lakes, with a fauna dominated by large branchiopods.

Momentary species richness correlated negatively with salinity in three of the five lakes: Wombah $r = -0.3700$, $n=27$, ns; unnamed lake $r = -0.6135$, $n=7$ ns; Bulla $r = -0.6245$, $n=31$ significant at $P < 0.01$; Mid

Blue $r = -0.5072$, $n=35$, significant at $P < 0.01$; North Blue $r = -0.4853$, $n=29$, significant at $P < 0.01$.

Benthos

Based on limited sampling, benthos of unvegetated offshore zone of the lakes was abundant but communities were simply structured (Table 5). Chironomids dominated, with ceratopogonids and large ostracods present, and an oligochaete in one lake. Biomass was by far the greatest in Lake Bulla.

Table 5. Benthic invertebrates of four of the lakes (numbers per m²).

| Lake | Wombah | Bulla | Mid Blue | Nth Blue |
|---|------------|-------------|------------|------------|
| TDS(gL ⁻¹) | 1.5 | 17.1 | 6 | 30.9 |
| Depth (m) | 1.5 | 2 – 2.4 | 1 – 1.55 | 0.2 |
| <i>Dero digitata</i> (Muller) | | | 22 ± 22 | |
| <i>Mytilocypris splendida</i> (Chapman) | | 2177 ± 166 | | |
| <i>Trigonocypris globulosa</i> De Deckker | | 1320 ± 219 | | |
| <i>Chironomus</i> sp. | 6600 ± 895 | | 100 ± 62 | |
| <i>Procladius</i> sp. | 310 ± 92 | 1364 ± 182 | 3250 ± 723 | 3720 ± 267 |
| <i>Tanytarsus</i> sp. | | 25655 ± 944 | | |
| Ceratopogonid larva | | 280 ± 55 | 250 ± 108 | 324 ± 83 |
| Total numbers (m ²) | 6910 ± 768 | 30796 ± 712 | 3622 ± 544 | 4044 ± 248 |
| Biomass (gm ²) | 9.1 ± 1.1 | 33.5 ± 0.9 | 4.5 ± 0.7 | 4.9 ± 0.4 |

Fish

A variety of fish occurred intermittently in the lakes. Bony Herring (*Nematalosa erebi*) and Spangled Perch (*Leiopotherapon unicolor*) were present in Lakes Bulla, Mid Blue and North Blue from at least mid 1998 to 2001 and Mosquito fish (*Gambusia holbrooki*) and Carp (*Cyprinus carpio*) were common in Lake Wombah over the same period. Other fish could have been present in these four lakes as the sampling technique was not designed to catch all species. No fish were seen in the unnamed lake.

There was a major fish kill in Lake Bulla in March 2001 at the same time as there was a bluegreen algal bloom. Carp in Lake Wombah died en masse in December 2001 as the salinity reached 30 gL⁻¹. Fish in the Blue lakes seemed to disappear slowly over 2001, after which piscivorous birds were rarely seen on these lakes.

Waterbirds

Forty-eight species of waterbirds were seen on the lakes; Wombah had 34 species, the unnamed lake 15, Bulla 43, Mid Blue 41 and North Blue 38. The most common species were Grey Teal (*Anas gracilis*) (particularly in Wombah), Pink-eared Duck (*Malacorhynchus membranaceus*) (particularly in Bulla) and Eurasian Coot (*Fulica atra*) (mainly in Bulla and North Blue). Other species present in appreciable numbers included: Hardhead (*Aythya australis*), the Australasian (*Tachybaptus novaehollandiae*) and Hoary-headed (*Poliiocephalus poliocephalus*) grebes, Black Swan (*Cygnus atratus*), Australasian Pelican (*Pelecanus conspicillatus*), Little Black Cormorant (*Phalacrocorax sulcirostris*), Little Pied Cormorant (*Phalacrocorax melanoleucos*), Silver Gull (*Larus novaehollandiae*), Black-winged Stilt (*Himantopus himantopus*) and Red necked Avocet (*Recurvirostra novaehollandiae*). Only Black Swans bred on the lakes during 1995-2004, mainly on the islands in Bulla.

Numbers fluctuated greatly between lakes and observations (Fig. 4 and Table 3). Mean numbers of birds on the lakes varied between 18.6 ha⁻¹ (North Blue) and 7.3 ha⁻¹ (Wombah) (assuming the lakes were on average three-quarters full) (Table 3). Variability was greatest in Bulla which had the greatest salinity range, and there was no correlation between species richness or bird numbers with salinity ($r = 0.1364$ and $r = -0.0839$ respectively).

DISCUSSION

Naturally salinised lakes in the eastern inland of

Australia have many distinctive features as itemised below, and with particular reference to the Rockwell-Wombah lakes.

1. Hydrological regime and geomorphic factors determine the presence of salt lakes and their salinity fluctuations (Hammer 1986; Williams 1998a; Timms 2006). When hydrologically closed, i.e. without overflow, bodies of water tend to accumulate salt. This explains the different salinity regimes in the Rockwell-Wombah lakes. Lake Wombah is the only one to be flushed by river flood water and then evaporation concentrates the isolated waters. Turbidity is greatest in Lake Wombah, due to the colloidal clays of Paroo River water (Timms 1999). Lake Bulla is generally the evaporative terminus of a blocked stream system flowing onto the Paroo floodplain, so it has accumulated the most salt and, associated with this, its waters are the clearest. Every few decades it overflows and presumably salt is lost so that it is only hypersaline as it nears dryness (similar to Lake Wyara on the other side of the Paroo floodplain – Timms 1998). Mid Blue and North Blue Lakes are intermediate settling basins; differences between them largely reflect the relative shallowness of North Blue Lake. The unnamed lake is the most intermittent of the five due to its relatively small catchment; salt accumulation consequently is quite modest.

2. For episodic saline lakes even 10 years is not long enough to encounter all the variations in environmental conditions. Judged from comparative rainfall averages at Rockwell during 1995-04 (366mm) compared to long term averages for the area (325 mm over 105 years at Boorara nearby – A. McGarth, pers. com; 302 mm over 75 years at Warroo nearby – M. Dunk, pers. com.), the Rockwell-Wombah lakes were probably full for longer than a long-term average during 1995-2004. On the other hand, if the lakes had been studied during the much drier 1910s to 1930s they would have held only a little water occasionally and would have been more saline on average. An even longer time frame would encompass climate change and quite different hydrological conditions (Bowler 1983; Pearson et al. 2003).

3. While there is a classic negative relationship between diversity and salinity in salt lakes (Hammer 1986; Timms 1993), other factors may mask it (Williams 1998b); in the case of these lakes only the littoral assemblage in Lake Bulla and the two Blue Lakes were significantly correlated with salinity. None of the other influencing factors considered by Williams (1998b) seems to be important in these lakes.

ECOLOGY OF EPISODIC SALINE LAKES




Figure 4. Variations in bird numbers in four of the lakes 1998-2003.

Indeed, the wide salinity tolerance of many species is in accord with that in the remainder of Australia so that it is only where lakes exceed 50 gL⁻¹ that salinity has a significant impact as a determinant of community structure (Williams et al. 1990; Williams 1998b).

4. Salt lake fauna is regionalised in Australia (Williams 1984; Timms 2007), though all areas share a dominance of crustaceans, particularly *Parartemia* spp., *Apocyclops dengizicus*, *Daphniopsis* spp., and a host of ostracods (*Diacypriis*, *Heterocypris*, *Reticypris*, *Platycypris*) including a few mytilocyprid genera (*Australocypris*, *Mytilocypris*, *Trigonocypris*) (Williams 1981; Pinder et al. 2002, 2004a, 2004b; Halse & McRae 2004; Timms 2007;). Lakes in more propitious climates have higher crustaceans such as amphipods (*Austrochiltonia* spp) and the isopod *Haloniscus searlei* which have no resistant stage and need at least dampness to survive dry periods, and calanoid copepods (*Calamoecia* spp.) and more ostracod genera (Williams 1984; Pinder et al. 2002). The eastern inland has a distinct regionalised fauna, though as more is being learnt about inland Western Australian fauna (Pinder et al. 2002, 2004a; Timms et al. 2006) many of the distinctive species of the east are being found in the west as well. Examples include recent discoveries of *Eocyclus parooensis*., *Daphniopsis queenslandensis*, *Moina baylyi*, *Celsinotum* spp., and *Trigonocypris globulosa* in inland Western Australia (Halse et al. 2000; Pinder et al. 2002, 2004; Timms et al. 2006). The eastern inland is still characterised by the unique presence of *Parartemia minuta*, a new *Daphnia* sp. in hyposaline waters, many halotolerant insects, and many monotypic genera whereas there has been multiple speciation in other areas, particularly in Western Australia (Geddes et al. 1981; Pinder et al. 2002, 2004a), and a greater relative importance of a variety of insects (mainly odonotans, corixids, notonectids, and coleopterans).

5. Not all lakes have a full complement of the regional fauna probably due to a mixture of local environmental factors, stochastic events and inadequate sampling. In the case of the Rockwell-Wombah lakes, species composition is typical for lakes in the middle Paroo with fluctuating salinity (Timms, 1993; 1999; Timms and Boulton, 2001). Notable absences include *Calamoecia canberra* (because waters are not turbid enough when fresh – Timms 2001b), and *Parartemia minuta* (because the lakes often have fish). Notable unusual occurrences include the snail *Coxiella gilesi* (absent in most other

Paroo lakes investigated), *Daphnia lumholtzi* (more common in these lakes than elsewhere in Paroo), and *Branchinella buchananensis* (this is the only known site in southwest Queensland). Some differences between the lakes can be explained in terms of various environmental factors. The presence of shrimps and yabbies in Lake Wombah when it is fresh-subsaline is explained by its episodic river connection. The importance of large branchiopods in the unnamed lake can be associated with the permanent absence of fish in it. Finally the longer species list of coleopterans in Mid Blue Lake (23 species cf 13 -17 in the other lakes) could be a reflection of this lake's greater littoral heterogeneity, among other factors.

6. Lakes that fill predictably each season have a shorter list of species than those fill episodically (areas being of the same magnitude). The Rockwell-Wombah lakes, together with other Paroo lakes and Lake Gregory in northwest Western Australia with episodically fluctuating salinities, have relatively long species lists (ca 70-100 species) (Table 2 and 3; Timms 1998; Halse et al. 1998; Timms & Boulton, 2001; Timms & McDougall 2004). By contrast, seasonally-filled salinas in southern Victoria (Geddes 1976) and southeast South Australia (Geddes & Brock 1978; De Deckker & Geddes 1980) have less than half this number (ca 25 -40 species). The explanation lies mainly in the significant periods of hyposaline conditions (these lakes have lower median salinities than most southern seasonally filled lakes), and also in the unpredictable and fluctuating conditions in the episodic lakes suiting various species at different times, compared with predictable and generally muted environmental conditions of the seasonal lakes allowing just one suite of tolerant species to persist throughout much of the season. Lack of salinity variation to maintain diversity may well be a major factor contributing to low species richness in secondarily salinised lakes (Hudson et al. 2003).

7. Habitat heterogeneity is another major factor influencing biodiversity in saline lakes (Timms 1998, 2001c; Williams 1998b), and needs to be considered in conjunction with salinity variation. Large salinas such as Lake Eyre are far more homogeneous than small salt lakes and have fewer species (Timms 1998). A Paroo example is a comparison of Lake Wyara (3400 ha) and nearby Lake Bulla (420 ha), both studied for 10 years, both exhibiting similar salinity fluctuations over this time and both studied by similar methods – with 34 species recorded in Lake Wyara, but 73 in Bulla (Timms 1998). This role for environmental heterogeneity as a species richness driver is confirmed

ECOLOGY OF EPISODIC SALINE LAKES

by the presence of 84 species in the Werewilka Inlet of Lake Wyara (Timms 2001c).

8. Distinctions between discrete zooplankton, littoral and benthic communities are blurred in most saline lakes, because most are shallow and many have rich macrophyte growth throughout the lake. The situation is not helped by the giant ostracods (*Australocypris*, *Mytilocypris*, *Trigonocypris* species) and the large branchiopods (*Parartemia* spp., *Eocyclus parooensis*, *Triops 'australiensis'* and halobiont species of *Branchinella*) which live and feed variously in all three habitats (Marchant and Williams 1977 on *Parartemia zietziana*; Timms 1981 on giant ostracods in L.Gnotuk; others from unpublished data, author). Many studies (e.g. Geddes 1976; Geddes & De Deckker 1980) therefore make little distinction between habitats, while others try, but with overlapping species lists (e.g. present work). Most undersample the benthos either by failing to dig the littoral net sufficiently into the sediments, or no or few specialised studies with quantitative devices such as Birge-Ekman grabs, or both. Thus while many benthic species seem to be recorded in so called littoral samples, indications of abundance are far too low (cf chironomids in Tables 4 and 5).

9. Hyposaline and mesosaline lakes attract waterbirds because of their rich resources (Kingsford et al. 1994; Kingsford & Porter 1994; Timms 1997). Even hypersaline lakes can support vast numbers of a limited diversity of birds, mainly large waders (Chapman & Lane 1997) and can be important breeding sites for some species (e.g. Banded Stilts—Burbridge & Fuller 1982). The Rockwell-Wombah lakes supported many thousands of birds belonging to 48 species. Numbers fluctuated widely (Fig. 4) due in part to varying food resources (Kingsford & Potter 1994; McDougall & Timms 2002), but these were not studied in these lakes. Mobile waterbirds are also influenced by events elsewhere in the inland. For example, the low numbers in the Rockwell-Wombah Lakes and also in nearby Lake Yumberarra during 2000 (Timms & McDougall 2004) probably marks their movement to the Lake Eyre Basin in response to even better conditions there (Roshier et al. 2002).

10. Saline lakes are usually productive, especially when hyposaline or mesosaline, but not when euhypersaline (Williams 1972; Hammer 1981a, 1981b; Timms 1983). In the Rockwell-Wombah lakes there are indications that Lake Bulla, the most saline lake is also the most productive (Table 3). However at any instant, some of the present lakes are unproductive

due to either the aftermath of a bluegreen algal bloom (as at Lake Bulla) or by filamentous algae overgrowing and killing macrophytes as seen in many summers at the Blue lakes.

ACKNOWLEDGEMENTS

This paper was first presented orally at a LIMPACS 'Salinity, Climate and Salinisation' conference at Mildura, Victoria in October, 2004. I sincerely thank John and Heather Buster, Robin and Rhonda Davis and family, James and Cheryl Hatch and Paul and Geraldine Tuite, variously present and former owners and managers of Rockwell and Wombah, for permission to work on their properties, for extraordinary generous hospitality, for assistance on occasions when I got hopelessly bogged, and to James and John for bulldozing tracks to make my passage easier between lakes. I appreciate the assistance over the years of many volunteer field assistants, including John Vosper and Sarah Wythes who came many times. I thank Olivier Rey-Lescure for drawing Figure 1, and Jane McRae, Adrian Pinder and Rus Shiel for identifications. Finally I am grateful to Jenny Davis and Lien Sim of Murdoch University, Perth, contributed with their helpful comments on the manuscript.

REFERENCES

- Bowler, J.M. (1983). Lunettes as indices of hydrologic change: a review of Australian evidence. *Proceedings of the Royal Society of Victoria* **93**: 147-168.
- Burbridge, A.A. and Fuller, P.J. (1982). Banded Stilts breeding at lake Barlee, Western Australia. *Emu* **97**: 212-216.
- Chapman, A. and Lane, J.A.K. (1997). Waterfowl usage of wetlands in the south-east arid interior of Western Australia 1992-93, *Emu* **97**: 51-59.
- Davis, J.A., McGuire, M., Halse, S.A., Hamilton, D., Horwitz, P., McComb, A.J., Froend, R.H., Lyons, M and Sim, L. (2003). What happens when you add salt: predicting impacts of secondary salinisation on shallow aquatic ecosystems by using an alternative-states model. *Australian Journal of Botany* **51**: 715-724.
- De Deckker, P. and Geddes, M.C. (1980). Seasonal fauna of ephemeral saline lakes near the Coorong Lagoon, South Australia. *Australian Journal of Marine and Freshwater Research* **31**: 677-699.
- De Deckker, P. and Williams, W.D. (1982). Chemical and biological features of Tasmanian salt lakes. *Australian Journal of Marine and Freshwater Research* **33**: 1127-1132.
- Geddes, M.C. (1976). Seasonal fauna of some ephemeral saline waters in western Victoria with particular reference to *Parartemia zietziana* Sayce (Crustacea: Anostraca). *Australian Journal of Marine and Freshwater Research* **27**: 1-22.

- Geddes, M.C. and Brock, M. (1978). Limnology of some lagoons in the southern Coorong. In (eds) 'The Southern Coorong and Lower Youngusband Peninsula of South Australia' (Eds. D.D. Gilbertson, and M.R. Foale) pp E47-8 (The Nature Conservation Society of South Australia Inc, Adelaide)
- Geddes, M.C., De Deckker, P., Williams, W.D., Morton, D.W. Topping, M. (1981). On the chemistry and biota of some saline lakes in Western Australia. *Hydrobiologia* **81/82**: 201-222.
- Halse, S.A., Shiel, R.J., Storey, A.W., Edward, D.H.D., Lansbury, I, Cale, D.J. and Harvey, M.S. (2000). Aquatic invertebrates and waterbirds of wetlands and rivers of the southern Carnarvon Basin, Western Australia. *Records of the Western Australian Museum Supplement*. **61**: 217-265.
- Halse, S.A. and McRae, J. (2004). New genera and species of 'giant' ostracods (Crustacea: Cyprididae) from Australia. *Hydrobiologia* **524**: 1-52.
- Halse, S.A., Ruprecht, J.K. and Pinder, A.M. (2003). Salinisation and prospects for biodiversity in rivers and wetlands of south-west Western Australia. *Australian Journal of Botany* **51**: 673-688.
- Hammer, U.T. (1981a). A comparative study of primary production and related factors in four saline lakes in Victoria, Australia. *Internationale Revue Gesamten Hydrobiologie* **66**: 701-743.
- Hammer, U.T., (1981b). Primary production in saline lakes. A review. *Hydrobiologia* **81**: 47-57.
- Hammer, U.T. (1986). 'Saline lake ecosystems of the world.' (Junk, Dordrecht).
- Hudson, P., Sheldon, F. and Costelloe, J. (2003). Aquatic macroinvertebrate biodiversity in the Western Lake Eyre Basin: The role of naturally fluctuating salinity. *Records of the South Australian Museum Monograph Series No. 7*: 135-144.
- Kingsford, R.T. and Porter, J.L. (1994). Waterbirds on an adjacent freshwater lake and salt lake in arid Australia. *Biological Conservation* **69**: 219-228.
- Kingsford, R.T., Bedward, M. and Porter, J.L. (1994). Waterbirds and Wetlands in northwestern New South Wales. Occasional Paper No. 19, NSW National Parks & Wildlife Service, Hurstville.
- Marchant, R., and Williams, W.D. (1977). Population dynamics and production of a brine shrimp *Parartemia zietziana* Sayce (Crustacea: Anostraca) in two salt lakes in western Victoria. *Australian Journal of Marine and Freshwater Research* **28**: 417-438.
- Pearson, S., Gayler, S, Hartig, K. and Timms, B. (2003). Ecosystem health in the Paroo: an arid frontier? In 'Proceedings of The Air Waters Places Transdisciplinary Conference on Ecosystem Health in Australia' (Ed, G. Albrecht) pp 252-264. (School of Environmental and Health Sciences, University of Newcastle)
- Pinder, A.M., Halse, S.A., Shiel, R.J., Cale, D.C. and McRae, J.M. (2002). Halophile aquatic invertebrates in the wheatbelt region of south-western Australia. *Verhandlungen Internationale Vereinigung Limnologie* **28**: 1687-1693.
- Pinder, A.M., Halse, S.A., McRae, J.M. and Shiel, R.J. (2004a). Occurrence of aquatic invertebrates of the Wheatbelt region of Western Australia in relation to salinity. *Hydrobiologia* **543**: 1-24. PROBLEM
- Pinder, A.M., Halse, S.A., McRae, J.M. and Shiel, R.J. (2004b). Aquatic invertebrate assemblages of wetlands and rivers in the Wheatbelt region of Western Australia. *Records of the Western Australian Museum Supplement* **67**: 7-37.
- Roshier, D.A., Robertson, A.I. and Kingsford, R.T. (2002). Responses of waterbirds to flooding in an arid region of Australia and implications for conservation. *Biological Conservation* **106**: 399-411.
- Timms, B.V. (1981). Animal communities in three Victorian lakes of differing salinity. *Hydrobiologia* **81**: 181-193.
- Timms, B.V. (1983). A study of benthic communities in some shallow saline lakes of western Victoria, Australia. *Hydrobiologia* **105**: 165-177.
- Timms, B.V. (1993). Saline lakes of the Paroo, inland New South Wales, Australia. *Hydrobiologia* **267**: 269-289.
- Timms, B.V. (1997). A comparison between saline and freshwater wetlands on Bloodwood Station, the Paroo, Australia, with special reference to their use by waterbirds. *International Journal of Salt Lake Research* **5**: 287-313.
- Timms, B.V. (1998). A study of Lake Wyara, an episodically filled saline lake in southwest Queensland, Australia. *International Journal of Salt Lake Research* **7**: 113-132.
- Timms, B.V. (1999). Local Runoff, Paroo Floods and Water Extraction Impacts on the Wetlands of Currawinya National Park. In 'A free-flowing river: the ecology of the Paroo River' (Ed R.T. Kingsford) pp. 51-66. (NSW National Parks and Wildlife Service, Sydney).
- Timms, B.V. (2001a). Wetlands of Currawinya National Park: Conservation and Management. In 'Research needs for managing a Changed Landscape in the Hungerford/Eulo Region – A Workshop held at Currawinya National Park 16th May 2001' (Eds. M. Page, C. Evenson and A. Whittington) pp.9-12. (University of Queensland, Gatton).
- Timms, B. V. (2001b). A new species of *Calamoecia* (Copepoda: Calanoida) from arid Australia, with comments on the calanoid copepods of the Paroo, northwestern Murray-Darling Basin. *Memoirs of the Queensland Museum* **46**: 783-790.
- Timms, B.V. (2001c). A study of the Werewilka Inlet of the saline Lake Wyara, Australia – a harbour of biodiversity for a sea of simplicity. *Hydrobiologia* **466**: 245-254.
- Timms, B.V. (2005). Salt lakes in Australia: present problems and prognosis for the future. *Hydrobiologia* **552**: 1-15.
- Timms, B.V. (2006). The geomorphology and hydrology of saline lakes of the middle Paroo, arid-zone Australia. *Proceedings of the Linnean Society of New South Wales* **127**: 157-174.
- Timms, B.V. (2007). The limnology of the saline lakes of central and eastern inland of Australia: A review with

ECOLOGY OF EPISODIC SALINE LAKES

- special reference to their biogeographical affinities. *Hydrobiologia* **576**: 27-37.
- Timms, B.V. and Boulton, A. (2001). Typology of arid-zone floodplain wetlands of the Paroo River, inland Australia and the influence of water regime, turbidity, and salinity on their aquatic invertebrate assemblages. *Archiv für Hydrobiologie* **153**: 1-27.
- Timms, B. V., Datson, B. and Coleman, M. (2006). The Wetlands of the Lake Carey Catchment, Northeast Goldfields of Western Australia, with special reference to large branchiopods. *Journal of the Royal Society of Western Australia* **89**: 175-183
- Timms, B.V. and McDougall, A. (2004). The limnology of Lake Yumberarra, an episodic arid zone wetland, with special reference to its use by waterbirds. *Wetlands (Australia)* **22**: 11-28.
- Williams, W.D. (1972). The uniqueness of salt lake ecosystems. In 'Productivity Problems of Freshwaters' Eds S. Kajak and A. Hillbricht-Ilkowska) pp. 349-361. (Polish Academy of Sciences, Warsaw).
- Williams, W.D. (1981). The limnology of saline lakes in Western Victoria. *Hydrobiologia* **82**: 233-259.
- Williams, W.D. (1984). Chemical and biological features of salt lakes on the Eyre Peninsula, South Australia, and an explanation of regional differences in the fauna of Australian salt lakes. *Verhandlungen Internationale Vereinigung Limnologie* **22**: 1208-1215.
- Williams, W.D. (1998a). 'Management of Inland Saline Waters. Guidelines of Lake Management Vol 6'. (International Lake Environment Committee United Nations Environment Programme. Kusatsu, Japan).
- Williams, W.D. (1998b). Salinity as a determinant of the structure of biological communities in salt lakes. *Hydrobiologia* **381**: 191-201.
- Williams, W.D. (2002). Environmental threats to salt lakes and the likely status of inland saline ecosystems in 2025. *Environmental Conservation* **29**: 154-167.
- Williams, W.D., Boulton, A.J. and Taafe, R.G. (1990). Salinity as a determinant of salt lake fauna: a question of scale. *Hydrobiologia* **197**: 257-266.